



2022 Water Quality Report

U.S. Army Corps of Engineers
Saint Louis District

Carlyle Lake Water Quality Conditions: 1971-2022



October 2023

Carlyle Lake Water Quality Conditions: 1971-2022

Prepared for

United States Army Corps of Engineers
Saint Louis District
1222 Spruce Street
Saint Louis, MO 63103-2833

Prepared by:

Ben Greeling
Environmental Specialist

EXECUTIVE SUMMARY

The United States Army Corps of Engineers (USACE) commitment to environmental compliance and protection of estuaries, rivers, lakes, and navigable waters arises from the national policy and directives expressed in Federal Statutes, Executive Orders, and internal regulations. These regulations were designed to minimize pollution, maximize recreation, protect aesthetics, preserve natural resources, and promote the comprehensive planning and use of water bodies to enhance the public interest rather than private gain. Therefore, USACE, in the design, construction, management, operation, and maintenance of its facilities, will exert leadership within existing authorities and appropriations in the nationwide effort to protect, enhance, and sustain the quality of the nation's resources. It is USACE's policy to comply with requirements of the Clean Water Act and not to degrade existing water quality conditions to the maximum extent that is practicable, consistent with project authorities, Federal legal and regulatory requirements, the public interest, and water control manuals.

The United States Army Corps of Engineers, Saint Louis District (CEMVS), implemented a water quality monitoring program during the 1970s to evaluate how its civil projects may be affecting water resources. Data collected from this effort serves as an invaluable tool for evaluating the significance of annual water quality measurements and tracking long-term trends. Water quality data is provided to the Missouri Department of Natural Resources and the Illinois Environmental Protection Agency to be used as a screening mechanism for the Missouri and Illinois Water Quality Report which is required every two years by the Clean Water Act Sections 303(d) and 305(b).

The National Water Quality Inventory Report to Congress (305(b) report) is the primary vehicle for informing law makers and the public about general water quality conditions in the United States. This document characterizes our water quality, identifies widespread water quality problems of national significance, and describes various programs implemented to restore and protect our waters. Currently the Illinois Environmental Protection Agency (IEPA, 2020) has listed Carlyle Lake as impaired for total phosphorous and mercury while the Kaskaskia River upstream from the Lake is impaired for fecal coliform, and mercury. In addition, the North Fork Kaskaskia River is impaired for phosphorus, Atrazine, and Terbufos. The lists of sources for these impairments are contaminated sediments, crop production, and unknown sources. The entire Kaskaskia watershed is impaired by the above parameters as well as many others.

Water quality sampling in 2022 revealed the following concerns at Carlyle Lake: dissolved oxygen, temperature, bacteria, total phosphorus, iron, pH, and chlorophyll a.

TABLE OF CONTENTS

INTRODUCTION.....	5
CARLYLE LAKE WQMP COVERAGE	6
Sample Location Summary Table	8
METHODS AND ANALYSIS: WATER QUALITY	9
Data Collection and Historical Reference Data	9
Statistical Summary and Comparison to Applicable Water Quality Standards.....	9
Quality Assurance	9
Water Quality Parameters and Criteria	10
Laboratory Methods and Water Quality Criteria Summary Table.....	15
RESULTS AND SUMMARY STATISTICS: WATER QUALITY	17
DISCUSSION: WATER QUALITY	33
MONITORING PROGRAM RECOMMENDATIONS	36
WORKS CITED	37
APPENDIX A: FIELD DATA	38
APPENDIX B: LABORATORY DATA.....	42

INTRODUCTION

The Carlyle Lake watershed encompasses approximately 1,663 square miles and includes all or portions of Bond, Clinton, Effingham, Fayette, Marion, Shelby, and Montgomery counties. The watershed includes the Kaskaskia River between Carlyle Lake Dam and Lake Shelbyville Dam and major tributaries of the Kaskaskia River, including: Big, Richland, Robinson, Becks, Ramsey, Old Hickory, and Hurricane Creeks (respectively) and the East Fork Kaskaskia River. Agriculture is the predominant land use within the watershed. Currently, 82% of the land is used for agricultural purposes. Of that 82%, 63% is cropland and 19% grassland. Since 1978, the number of farms has decreased by 25% and the acreage tilled has decreased by only 6%. Corn and soybeans are important to the region, but producers also grow 25% of the entire state's crop of wheat. Livestock production, including dairy, swine, poultry, and beef cattle is a significant industry, especially in Clinton, Randolph, and Washington Counties.

Carlyle Lake is located in south central Illinois at river mile 94.2 on the Kaskaskia River, upstream from its confluence with the Mississippi River and about one-half mile upstream from the town of Carlyle, Illinois. Carlyle is located in Clinton County, approximately 50 miles east of St. Louis, Missouri. Carlyle Lake is the largest man-made lake in the state and is approximately 12 miles long and 1-3 miles wide and has approximately 24,710 acres of water surface at summer pool elevation 445.0 feet NGVD (National Geodetic Vertical Datum). There are 88 miles of shoreline and approximately 12,800 acres of public land associated with the project. The lake is situated in gently rolling land with alluvial valleys with moderately low relief. The lake provides outdoor recreation opportunities for over 2.5 million visitors annually, which generates over \$80 million in visitor spending within 30- miles of the Lake. There are 41 recreation areas that include: 424 picnic sites, 726 campsites, 670 marina slips, 24 boat ramps, and 25 miles of hiking trails. The CEMVS manages and operates two large reservoirs on the Kaskaskia River, Lake Shelbyville and Carlyle Lake, as well as the 36 mile long navigable channel and lock and dam at the Kaskaskia River Project.

Carlyle Lake is managed and operated by the CEMVS for the authorized purposes of flood risk management, navigation, water supply, water quality, fish and wildlife conservation, and recreation. The lake serves as a heavy recreational usage lake. The land surrounding the lake is used predominately for agriculture. Surrounding communities have existing industrial/commercial operations and residents which discharge wastewater into municipal wastewater treatment plants that ultimately discharge treated water into the Kaskaskia River basin. Agricultural runoff and municipal wastewater treatment facilities are the primary potential source of pollution into the Carlyle Lake watershed. Additional sources are marinas, recreational watercraft discharges and wildlife fecal material runoff.

Water quality is of paramount importance for sustaining ecological integrity and services provided by the Kaskaskia River and Carlyle Lake. Water quality is influenced by a range of both point and nonpoint pollution sources, which may include natural

processes, industrial and municipal effluents, and surface runoff from agricultural arenas.

The USACE has implemented a Water Quality Management Plan (WQMP) as part of the operation and maintenance activities associated with managing USACEs' civil works projects throughout the District which includes, among other reservoirs and rivers, the Kaskaskia River and Carlyle Lake. The WQMP addresses surface water quality management issues and adheres to the guidance and requirements specified by Clean Water Act (CWA), as well as the self-imposed Engineering Regulation (ER) 1110-2-8154, "Water Quality and Environmental Management for USACE Civil Works Projects" (USACE, 2018). Water quality monitoring is implemented to fulfill five primary objectives that drive the CEMVS WQMP:

- 1) Establish baseline conditions, identify significant water quality trends, and document problems and accomplishments.
- 2) Ensure that surface water quality, as affected by CEMVS projects, is suitable for project purposes, existing water uses, public health and safety, and in compliance with applicable state and federal water quality standards.
- 3) Provide support to water control, project operations, and navigation for regulations and modifications.
- 4) Investigate special problems, design and implement modifications, and improve water management procedures
- 5) Establish and maintain strong working partnerships and collaborations with appropriate entities within and outside USACE regarding water quality.

This report is intended to document and assess water quality conditions occurring at Carlyle Lake. The report describes conditions observed in 2022, as well as baseline data collected from 1971-2021. Data are available upon request.

CARLYLE LAKE WQMP COVERAGE

The WQMP for Carlyle Lake includes water samples taken at the following locations: major tributaries (CAR-13 and CAR-12), main body of the lake (CAR-4, CAR-2, and the marinas), and just downstream of the dam (CAR-1). See figure 1 and Table 1 for a site map and site coordinates.




Figure 1. Water Quality (WQ) Sampling Locations in 2022 at Carlyle Lake

Sample Location Summary Table

Table 1: Sample Location Summary and Geographic Location (NAD 1983)

Sample Location Type	Abbreviation	Site Name	Latitude	Longitude
Major Tributary	TRIB	CAR-13	38.868961	-89.159605
	TRIB	CAR-12	38.868961	-89.193475
Main Reservoir Surface	RS	CAR-2	38.619492	-89.352747
	RS	CAR-4	38.740632	-89.267266
	RS	CAR-BL	38.693092	-89.234040
	RS	CAR-DW	38.627955	-89.358246
	RS	CAR-KP	38.736930	-89.273674
	RS	CAR-CSA	38.642647	-89.336805
Reservoir Benthic	RB	CAR-2-10	38.619492	-89.352747
Tail Race (below dam)	TR	CAR-1	38.616240	-89.355828

Samples at Marinas are not always taken in the exact same location. *BL=Boulder Marina, DW=Dam West Marina, KP=Keyesport Marina, CSA=Carlyle Sailing Association.*

METHODS AND ANALYSIS: WATER QUALITY

Data Collection and Historical Reference Data

During 2022, water quality samples were collected and analyzed for 10 locations during four separate sampling events (n=40; Table 1). One duplicate sample was also collected during each sampling event for quality control purposes. Samples were collected from the upper one meter of the water column, preserved, and transported to the Applied Research and Development Laboratory (ARDL) in Mount Vernon, Illinois for analysis.

For the purpose of this report, historical reference data refers to water quality data collected during the previous years ranging as far back as 1971 (parameter dependent) at Carlyle Lake. Historical reference data are intended to represent the current condition of Carlyle Lake.

Statistical Summary and Comparison to Applicable Water Quality Standards

Statistical analyses were performed on water quality monitoring data collected for 10 locations, and classified as TRIB (n= 2), RS (n=6, RB (n=1), and TR (n=1). For comparison, statistical analyses were also performed on historical water quality monitoring data and, although some sampling locations may have been removed, they were classified in the same manner. Descriptive statistics were calculated to describe central tendencies and boxplots created to illustrate comparisons between groups. Monitoring results were compared to applicable water quality standard criteria established by the appropriate state agencies pursuant to the Federal Clean Water Act. If state water quality standard criteria were not available, recommended criteria from the literature were considered.

Seasonal data are classified as: Winter (December 01 - March 14), Spring (March 15 – May 31), Summer (June 1 – September 15), Fall (September 16 – November 30).

Quality Assurance

The United States Army Corps of Engineers, Saint Louis District quality assurance procedures considers two primary focus areas: (1) those that involve laboratory analysis of samples, and (2) those concerning the collection and processing of the water samples in the field.

Since 2012, ARDL has analyzed water quality samples for CEMVS. Their quality assurance program includes the use of quality control charts, check standards, field and in-house matrix spikes, laboratory blanks and performance evaluation samples. In addition, one blind duplicate sample is submitted for at least every 20 samples, or, in this case, every sampling event (one event/day at Carlyle Lake has 6 lab samples and one duplicate).

Internal checks are also used for field sampling. This includes adherence to operating procedures for data collection and periodic evaluation of sampling personnel. Field sampling equipment and multimeters are calibrated/serviced in accordance with factory recommendations.

Water Quality Parameters and Criteria

Parameters used to characterize water quality have been generally accepted criteria for assessing aquatic life and human health include:

Temperature (Temp) is important because it controls several aspects of water quality. Colder water holds more dissolved oxygen which is required by aquatic organisms. Plants grow more rapidly and use more oxygen in warmer water. Decomposition of organic matter which uses oxygen is accelerated in warmer water. Temperature can also determine the availability of toxic compounds such as ammonia. Since aquatic organisms are cold blooded, water temperature regulates their metabolism and ability to survive. The number and kinds of organisms that are found in streams or lakes is directly related to temperature. Certain organisms require a specific temperature range, such as Salmonids, which require water temperatures below 20°C. Water temperature criteria for warm water bodies in Illinois is within 2.8°C of the seasonal norm.

Dissolved Oxygen (DO) refers to the measurement of free oxygen molecules (O_2) that are not bonded to any other elements; thus, oxygen bonded in water (H_2O) would not be considered in a measurement of dissolved oxygen. Oxygen is dissolved in surface waters through interactions with the atmosphere and as a waste product of photosynthesis ($CO_2 + H_2O \rightarrow CH_2O + O_2$) from phytoplankton and aquatic vegetation. Additional factors influencing DO include temperature, pressure, and salinity.

Dissolved oxygen is required for most aquatic life including fish, invertebrates, bacteria, and plants. Fish and invertebrates utilize DO for respiration through gills and cutaneous breathing, and plants require dissolved oxygen for respiration when photosynthesis is not possible. Smaller microbes and bacteria utilize DO for decomposition of organic materials, a process essential for nutrient cycling. Bottom feeders such as worms and mussels can persist when DO is ≥ 1 mg/L, while most inland fish species require a minimum DO of 4mg/L. The DO water quality criteria for Illinois is ≥ 5 mg/L.




Figure 1: Dissolved oxygen (O_2) vs oxygen bonded in water (H_2O).

Potential of Hydrogen (pH) is a measure of how acidic or basic water is. Potential of Hydrogen is reported on a logarithmic scale ranging from 0 – 14, with 7.0 being neutral. As pH increases from 7.0, water increases in alkalinity, whereas a decrease from 7.0 indicates an increase in acidity. Since pH is measured on a logarithmic scale, every

one-unit change in pH indicates a 10-fold change in acidity; thus, a pH of 6.0 is ten times more acidic than a pH of 7.0 and a pH of 4.0 would be one-thousand times more than a pH of 7.0.

The pH of water varies considerably beyond the local level. Natural variation in bedrock and soil composition through which water moves has been reported as one of the most influential factors. Additional factors include decomposition of organic materials, acidity of local precipitation, discharge of effluents and chemicals, and mining operations.

Most freshwater streams and rivers have a natural pH ranging from 6 to 8. As pH approaches 5 (acidic), less tolerant fish and aquatic invertebrate assemblages may be extirpated, and a pH below 4.5 would be without most desired aquatic life. Conversely, when pH exceeds 9.5 (alkaline), aquatic fish and invertebrate begins to rapidly decrease and beyond 10, fish become extirpated. The pH water quality criteria for Illinois ranges from 6.5 – 9.0.

Conductivity is a measure of water's ability to conduct electrical current. In its purist form, water has a *near* neutral charge, indicating that it is an inefficient conductor of electrical current. Thus, the ability to carry electrical current is driven by water soluble ions (atoms and molecules with a charge) such as salts and other inorganic materials. Conductivity is also influenced by water temperature; as temperature increases, conductivity increases. For this reason, conductivity is commonly reported as Specific Conductivity (SpCond), which is the measurement of conductivity at 25 degrees Celsius.

Conductivity in streams and rivers is affected by the geology of the area. Streams running through granite tend to have lower conductivity due to granite being composed of inert material; materials that do not ionize or dissolve into ionic compounds in water. Conversely, streams that run through areas of limestone or clay soils tend to have higher conductivity readings because of the presence of materials that ionize. Conductivity is useful as a general measure of water quality. A stream tends to have a relatively constant range of conductivity that, once established, can be used as a baseline. Significant changes, either increases or decreases, might indicate a source of pollution has been introduced into the water. The pollution source could be a treatment plant, which raises the conductivity, or an oil spill, which would lower the conductivity. In general, there are no water quality criteria for SpCond. The District threshold of 500 $\mu\text{S}/\text{cm}$ is a rule of thumb value that is often associated with some form of biological impairment.

Oxidation Reduction Potential (ORP) is a measurement of the net status of all the oxidation and reduction reactions in a given water sample. Oxidation involves an exchange of electrons between 2 atoms. The atom that loses an electron is oxidized and the one that gains an electron is reduced. Oxidation reduction potential sensors measure the electrochemical potential between the solution and a reference electrode. Readings are expressed in millivolts. Positive readings indicate increased oxidizing potential and negative readings increased reduction. Oxidation reduction potential

values are used much like pH values to determine water quality. While pH readings characterize the state of a system relative to the receiving or donating hydrogen ions (base or acid), ORP readings characterize the relative state of losing or gaining electrons. Generally, ORP readings above 400mV are harmful to aquatic life; however, ORP is a non-specific measurement, which reflects a combination of effects of all the dissolved materials in the water. Therefore, the measurement of ORP in relatively clean water has only limited utility unless a predominant redox-active material is known to be present.

Total Suspended Solids (TSS) concentrations, which cause the photosynthetic activity to be reduced by more than 10% from the seasonably established norm, can have a detrimental effect on aquatic life. Soil particles, organic material, and other debris comprise suspended solids in the water column. **Turbidity (FNU)** measurements are inverse to suspended solid measurements. As TSS increases, the FNU or water transparency decreases. Total suspended solids can be an important indicator of the type and degree of FNU. Total Suspended Solids measurements represent a combination of **Volatile Suspended Solids (VSS)**, which consist of organic material, and **Nonvolatile Suspended Solids (NVSS)**, which is comprised of inorganic mineral particles in the water. In order to more accurately determine the types and amounts of suspended solids, VSS are analyzed. Volatile suspended solid concentration represents the organic portion of the total suspended solids. Organic material often includes plankton, and additional plant and animal debris present in water. Total VSS indicates the presence of organics in suspension; and, therefore, show additional demand levels of oxygen. The Illinois Environmental Protection Agency suggests that generally NVSS above 15 mg/L could highly impair recreational lake use while NVSS of 3 to 7 mg/L may cause slight impairment (Hudson, 1998). Illinois does not currently have standard criteria for TSS, NVSS or VSS.

Total Organic Carbon (TOC) is a measure of the amount of organic carbon in a water body. In addition to natural organic substances, TOC includes insecticides and herbicides, as well as domestic and industrial waste. Industrial waste effluent may include carbon-containing compounds with various toxicity levels. Further, a high organic content means an increase in the growth of microorganisms which contribute to the depletion of oxygen supplies.

Currently, there are no state or federal water quality standard criteria set for TOC. Because carbon occurs naturally, its concentration varies based on physical and chemical attributes in a watershed; thus, this study relies on historical reference conditions to identify unfavorable conditions.

Metals Iron (TFe) and Manganese (TMn) (T=total) are nutrients for both plants and animals. Living organisms require trace amounts of metals. However, excessive amounts can be harmful to the organism. Heavy metals exist in surface waters in three forms, colloidal, particulate, and dissolved. Water chemistry determines the rate of adsorption and desorption of metals to and from sediment. Metals are desorbed from the sediment if the water experiences increases in salinity, decreases in redox potential,

or decreases in pH. Metals in surface waters can be from natural or human sources. Metal levels in surface water may pose a health risk to humans and the environment.

Pesticides are commonly used throughout much of the agricultural landscape that the Kaskaskia River flows. This study considers one insecticide and seven herbicides. Atrazine and Alachlor herbicides are commonly used agricultural chemicals which can be readily transported by rainfall runoff. Both compounds are suspected of causing cancer; and therefore, were monitored for the protection of human and aquatic health. Herbicides which are pesticides used to kill vegetation are the most widely used and sampled. Two of the most widely used herbicides are Atrazine and Alachlor. Atrazine is a preemergence or postemergence herbicide use to control broadleaf weeds and annual grasses. Atrazine is most commonly detected in ground and surface water due to its wide use, and its ability to persist in soil and move in water. Alachlor is a Restricted Use Pesticide (RUP) due to the potential to contaminate groundwater. The water quality standards for the pesticides sampled are located in Table 2.

Nitrogen occurs naturally in water through several forms including nitrogen (N₂), nitrite (NO₂-N), nitrate (NO₃-N), ammonia (NH₃), and ammonium (NH₄). Nitrates are the most commonly reported form of nitrogen and may have a meaningful influence on a water body's trophic status. Algae and other plants use NO₃-N as a food source, thus excess levels of NO₃-N can promote increases in algae production and hypereutrophic conditions.

In general, NO₃-N does not have a *direct* effect on fish or aquatic insects. Illinois has set criteria standards for NO₃-N to 10 mg/L to accommodate safe drinking waters for human and livestock; however, this threshold likely exceeds the concentration that is appropriate for assessing ecosystem health.

Total Ammonia Nitrogen (TAN) includes NH₃ and NH₄. Total ammonia nitrogen is a colorless gas with a strong pungent odor. Ammonia occurs naturally and is a biological requirement for aquatic life, however elevated concentrations can be toxic to freshwater organisms. Unnatural sources of ammonia include, accidental releases of ammonia rich fertilizer, effluent from sewage treatment plants, improper disposal of ammonia products, and livestock waste.

Toxic concentrations for freshwater organisms range from 0.53 – 22.8 mg/L and are strongly dependent on both pH and temperature. In general, an increase in pH and/or temperature corresponds with an increase in toxicity. Additional information in regard to the relationship between pH, temperature, and ammonia, as it relates to toxicity, can be reviewed in Aquatic Life Ambient Water Quality Criteria for Ammonia – Freshwater (USEPA 2013).

Total Phosphorus (TP) is analyzed as phosphorus and has been monitored due to the potential for uptake by nuisance algae. Levels of phosphate can indicate the potential for rapid growth of algae (algae bloom) which can cause serious oxygen depletion during the algae decay process. Phosphorous is typically the limiting nutrient in a water

body; therefore, any addition of phosphorous to the ecosystem stimulates the growth of plants and algae. Phosphorous is delivered to lakes and streams by way of runoff from agricultural fields and urban environments. Other sources of phosphorous are anaerobic decomposition of organic matter, leaking sewer systems, and point source pollution. The general standard for phosphorous in lake water is 0.05 mg/L. Dissolved phosphorous, also called **Orthophosphate (PO₄-P)** is generally found in much smaller concentrations than total phosphorous and is readily available for algal uptake. Orthophosphate concentrations in a water body vary widely over short periods of time as plants take it up and release it.

Chlorophyll a (CHL_a) is a measure of the number of algae growing in a waterbody, and therefore can be used to classify trophic status. Although algae are a natural part of freshwater ecosystems, too many algae can cause aesthetic problems such as green scums and bad odors and can result in decreased levels of DO.

Pheophytin a (PHEO_a) is a natural degradation product or digestion of CHL_a. The ratio of PHEO_a to CHL_a can provide an indication of the decline or growth in eukaryotic algae and cyanobacteria populations.

Trophic Status is determined using a modified **Trophic State Index (TSI)**, as described by Carlson (1977). Trophic State Index is calculated from Secchi-depth transparency, total phosphorus, and chlorophyll-a measurements. Values for these three parameters are converted to an index number ranging from 0-100 according to the following equations:

$$\begin{aligned} \text{TSI (Secchi Depth)} &= 10(6 - (\ln \text{SD}/\ln 2)) \\ \text{TSI (Chlorophyll-a)} &= \text{TSI(Chl)} = 10(6 - ((2.04 - 0.68 \ln \text{Chl})/\ln 2)) \\ \text{TSI (Total Phosphorus)} &= \text{TSI(TP)} = 10(6 - (\ln (48/\text{TP})/\ln 2)) \end{aligned}$$

where *ln* indicates the Natural Logarithm

A TSI average value, calculated as the average of the three individually determined TSI metrics, is used as an overall indicator of a water body's trophic state. The relationship between TSI and trophic condition is defined as follows:

TSI	Trophic Condition
0-40	Oligotrophic
40-50	Mesotrophic
50-70	Eutrophic
70-100	Hypereutrophic

Laboratory Methods and Water Quality Criteria Summary Table

Table 2: Metrics, Methods, and Water Quality Criteria Used for Evaluating Water Quality


Metric	Abbreviation	Analysis Method	Water Quality Criteria	Source
Alachlor		EPA Method 8270C	< 2µg/L PWS or <1100 µg/L: aquatic life	Illinois EPA
Ammonia Nitrogen	NH ₃	EPA Method 350.1	<15 mg/L	Illinois EPA
Atrazine	Atrazine	EPA Method 8270C	9 µg/L: Chronic or 82 µg/L: Acute or 3 µg/L DWS	Illinois EPA
Bacteria: E. Coliform	E Col	EPA Method 1604	< 235 E. Col per 100/mL for single sample	Illinois EPA
Chlorophyll a	Chl_a	SM Method 10200H	< 25 mg/m ³ (Eutrophic Upper Limit)	Carlson 1977
Chlorpyrifos		EPA Method 8270C	< .11 µg/L: aquatic life	Illinois EPA
Cyanazine		EPA Method 8270C	< 30 µg/L: chronic or < 370 ug/L acute (aquatic life)	Illinois EPA
Depth	Depth	Multiparameter Meter	Measurements reported at ~1 meter	-----
Dissolved Oxygen	DO	Multiparameter Meter	Greater than 5.0mg/L	Illinois EPA
Metolachlor		EPA Method 8270C	30.4 µg/L: Chronic or 380 µg/L: Acute	Illinois EPA
Metribuzin		EPA Method 8270C	8.4 mg/L: aquatic life or 8.3 mg/L: human health	Illinois EPA
Nitrate as Nitrogen	NO ₃	Green Method	< 10 mg/L	Illinois EPA
Non-Volatile Suspended Solids	NVSS	TSS - VSS	-----	-----
Orthophosphate	Ortho	EPA Method 365.2	-----	-----
Pendmethalin		EPA Method 8270C	< 30 µg/L: chronic or < 350 µg/L acute (aquatic life)	Illinois EPA
Pheophytin a	Phpy_a	SM Method 10200H	-----	-----
Potential of Hydrogen	pH	Multiparameter Meter	Range: 6.5 – 9.0pH	Illinois EPA
Specific Conductivity	SpCond	Multiparameter Meter	500 µS/cm	-----
Temperature	Temp	Multiparameter Meter	Less than rise of 2.8°C above normal seasonal temperature	Illinois EPA
Total Dissolved Solids	TDS	Multiparameter Meter	< 500 mg/L	Illinois EPA
Total Manganese	TMn	EPA Method 6010C	< 1 mg/L	Illinois EPA

Metric	Abbreviation	Analysis Method	Water Quality Criteria	Source
Total Organic Carbon	TOC	EPA Method 415.1	-----	-----
Total Iron	TFe	EPA Method 6010C	< 1 mg/L	Illinois EPA
Total Phosphorus	TP	EPA Method 365.2	Less than 0.05 mg/L	Illinois EPA
Total Suspended Solids	TSS	EPA Method 160.2	-----	Illinois EPA
Trifluralin		EPA Method 8270C	< 1.1 µg/L: chronic or < 26 µg/L acute (aquatic life)	Illinois EPA
Turbidity	Turb	Multiparameter Meter	-----	-----
Volatile Suspended Solids	VSS	EPA Method 160.4	-----	-----


**1 mg/L is equivalent to 1 drop in two bathtubs and 1 ug/L is equivalent to 1 drop in an Olympic size swimming pool. PWS is public water supply. DWS is drinking water standard.*

RESULTS AND SUMMARY STATISTICS: WATER QUALITY


Oxidation Reduction Potential: 1986–2021




Oxidation Reduction Potential: 2022



Specific Conductivity: 1971–2021







Specific Conductivity: 2022



Historical Reference 1971-2021					2022		
	Location	Mean	Median	n	Mean	Median	n
ORP	RB	228.07	231.0	134	----	----	----
	RS	242.74	235.0	277	150.35	161.3	16
	TR	283.03	296.0	132	208.17	172.1	3
	TRIB	278.85	286.5	172	185.70	164.5	5
SpCond	RB	405.18	411.0	239	----	----	----
	RS	406.99	411.0	637	341.77	334.8	22
	TR	397.38	395.0	248	357.68	361.4	4
	TRIB	518.89	495.0	605	392.44	414.7	7

*This report does not acknowledge a water quality criteria for SpCond or ORP.




* Red line placed at the 5 mg/L level for DO.


	Historical Reference 1971-2021				2022		
	Location	Mean	Median	n	Mean	Median	n
DO	RS	9.02	8.89	652	7.48	6.64	24
	TR	9.40	9.01	248	7.99	8.10	4
	TRIB	8.63	8.50	601	9.22	9.16	8
Temp	RS	19.13	21.00	661	24.56	24.76	24
	TR	18.93	20.85	254	23.95	24.58	4
	TRIB	15.74	17.71	618	24.16	23.29	8

* During the four sampling events in 2022 surface water DO was measured below the standard at CAR-4 and BL-MAR in June and at BL-MAR, DW-MAR, and KP-MAR in September. In 2022 temperature was recorded above the standard (rise of 2.8° C above the natural temperatures) during the spring. The historical seasonal mean temperature by class was used as the natural temperature.

Carlyle Lake Tailwater Water Quality 2022





**Data recorded by multi-parameter sonde at tail race in 2022. 15 minute data shown. DO was recorded below the standard of 5 mg/L several times throughout the year usually during low flow periods.*





*Red lines indicate the upper and lower water quality criteria standards (9 and 6.5).

	Historical Reference 1971-2021				2022		
	Location	Mean	Median	n	Mean	Median	n
pH	RS	7.99	8.00	649	8.22	8.22	24
	TR	7.88	7.90	250	8.04	8.01	4
	TRIB	7.56	7.60	603	7.94	8.02	8

*The pH standard was exceeded in May 2022 with readings greater than 9 at CAR-2, CAR-CSA, and CAR-KP.




*Red line indicates the water quality standard (10 mg/L).




Historical Reference 1971-2021					2022		
	Location	Mean	Median	n	Mean	Median	n
NO ₃ -N	RS	1.10	0.73	613	0.17	0.13	8
	TR	0.90	0.62	250	0.32	0.20	4
	TRIB	1.64	0.90	605	1.02	0.81	8
NH ₃ N	RS	0.08	0.05	508	0.07	0.08	8
	TR	0.14	0.09	216	0.08	0.07	4
	TRIB	0.12	0.07	424	0.07	0.08	8

*All observations of nitrate and ammonia nitrogen were within the water quality standard.


Orthophosphate: 1971–2021



Total Phosphorus: 1971–2021




Total Phosphorus: 2022



*Red line indicates the water quality standard of 0.05 mg/L.

Historical Reference 1971-2021					2022		
	Location	Mean	Median	n	Mean	Median	n
PO4	RS	0.08	0.06	600	----	----	----
	TR	0.10	0.07	244	----	----	----
	TRIB	0.07	0.05	598	----	----	----
TP	RS	0.22	0.16	620	0.26	0.27	8
	TR	0.23	0.19	249	0.32	0.35	4
	TRIB	0.24	0.19	616	0.20	0.16	8

*Total phosphorus exceeded the standard of 0.05 mg/L for all locations in 2022. This study does not acknowledge a standard for orthophosphate.




*Red line indicates the reference water quality standard of 25 mg/cm³. See Carlson 1977.


	Historical Reference 1984-2021				2022		
	Location	Mean	Median	n	Mean	Median	n
Chl_a	RS	30.31	21.40	282	47.90	44.55	8
	TR	32.30	20.50	5	30.10	24.90	4
	TRIB	23.62	11.90	68	46.71	49.30	8

*The reference standard for chlorophyll-a of 25mg/cm³ was exceeded at the most sites throughout 2022. This study does not acknowledge a standard for pheophytin.





Secchi Depth: 2022



2022 Carlson Trophic State Index




<40 = Oligotrophic __ 40-50 = Mesotrophic __ 50-70 = Eutrophic __ >70 Hypereutrophic




Historical Reference 2018-2021					2022		
	Location	Mean	Median	n	Mean	Median	n
TDS	RB	212.17	215.50	12	----	----	----
	RS	215.05	210.50	78	215.13	229.50	24
	TR	216.81	220.00	16	226.00	240.50	4
	TRIB	287.16	281.00	31	303.63	311.50	8
FNU	RB	RB	24.26	19.54	----	----	----
	RS	RS	39.28	25.04	36.85	21.99	24
	TR	TR	18.34	15.49	16.21	15.62	4
	TRIB	TRIB	66.40	45.74	27.85	26.11	8

* All observations of TDS were within the referenced water quality standard.

Total Suspended Solids: 1974–2021




Total Suspended Solids: 2022




Historical Reference 1974-2021					2022		
	Location	Mean	Median	n	Mean	Median	n
TSS	RB	41.59	29.00	229	29.95	31.15	4
	RS	34.66	25.00	521	27.71	26.30	8
	TR	25.16	21.00	218	22.50	18.75	4
	TRIB	105.86	62.25	436	57.66	54.45	8

**The mean total suspended solids data measured in 2022 were lower overall when compared to the historical data. There is no numeric standard for TSS.*

Total Organic Carbon: 1984–2021





Total Organic Carbon: 2022





Historical Reference 1984-2021					2022		
	Location	Mean	Median	n	Mean	Median	n
TOC	RS	6.51	5.60	318	5.09	5.04	8
	TR	6.58	5.50	150	4.71	4.68	4
	TRIB	6.03	5.00	209	4.01	3.80	8

**This study does not recognize a water quality criteria for TOC.*




*Red line indicates the water quality standard of 1 mg/L.



*Red line indicates the water quality standard of 1 mg/L.

	Historical Reference 1984-2021				2022		
	Location	Mean	Median	n	Mean	Median	n
TFe	RB	1.22	0.83	182	1.10	1.21	4
	RS	1.44	0.81	170	----	----	----
	TR	0.87	0.65	151	0.78	0.72	4
	TRIB	3.82	2.31	68	----	----	----
TMn	RB	0.23	0.18	182	0.18	0.18	4
	RS	0.16	0.14	160	----	----	----
	TR	0.19	0.15	149	0.17	0.16	4
	TRIB	0.33	0.26	61	----	----	----

*In 2022 iron exceeded the standard of 1 mg/L near the lake bottom in front of the dam in May, June, and August, and in the discharge once. Manganese did not exceed the criterion.




*Red line indicates the standard of 3 ug/L.


Historical Reference 1996-2021					2022		
	Location	Mean	Median	n	Mean	Median	n
Atrazine	RS	1.08	0.51	128	0.59	0.59	2
	TR	0.70	0.53	64	0.33	0.33	1
	TRIB	1.66	0.55	123	0.78	0.78	2

*Atrazine did not exceed the DWS criterion of 3 ug/L in 2022.

Metolachlor: 2007–2021




Metolachlor: 2022




Historical Reference 2007-2021					2022		
	Location	Mean	Median	n	Mean	Median	n
Metolachlor	RS	1.22	0.43	90	0.90	0.90	2
	TR	0.84	0.43	45	0.53	0.53	1
	TRIB	1.27	0.40	85	2.33	2.33	2

*Metolachlor did not exceed water quality criteria in 2022.

Metribuzin: 1998–2021




Metribuzin: 2022




Historical Reference 1998-2021					2022		
	Location	Mean	Median	n	Mean	Median	n
Metribuzin	RS	0.33	0.21	101	0.21	0.21	2
	TR	0.22	0.21	45	0.22	0.22	1
	TRIB	0.34	0.22	93	0.42	0.42	2

*Metribuzin did not exceed water quality criteria in 2022.

Surface Water E. Coli: 1996–2021



Surface Water E. Coli: 2022



*Red line indicates the water quality standard of 235 col per 100 mL.

Historical Reference 2001-2021					2022			
	Location	Mean	Median	n	Location	Mean	Median	n
E col	1	15.25	16.50	4	12	512.50	550.00	4
	2	29.00	8.00	13	2	139.75	78.00	4
	4	494.14	6.50	14	13	656.25	400.00	4
	BL	176.00	33.50	48	BL	141.50	140.00	4
	CSA	178.71	150.00	7	CSA	167.25	162.50	4
	DW	235.68	45.00	50	DW	290.75	137.50	4
	KP	175.02	71.00	49	KP	311.75	262.50	4

*Marina bacteria levels exceeded the standard throughout the lake and in the tributaries during the four sampling events in 2022.

2022 Swimming Beach/Marina Bacteria Levels (E. Coli / 100mL)

Month/week	McNair	Keyesport	Dam West	Harbor Light	Coles Creek	Boulder Marina	Carlyle Sailing Association Marina	West Access Marina	Trade Winds Marina
May/ week 1	55	43	57	112	77	43	25	75	67
2	34	54	41	156	87	65	43	154	133
3	57	78	45	154	89	120	76	88	133
4	76	67	86	133	122	75	43	122	96
June/week 1	75	56	86	186	146	122	78	122	155
2	66	45	57	166	122	95	53	75	164
3	40	20	20	170	80	60	69	45	90
4	86	54	65	120	1602	100	43	22	120
5	65	48	58	188	132	112	54	142	154
July/ week 1	77	123	65	200	148	112	58	147	165
2	76	43	78	168	142	141	76	180	160
3	58	96	112	200	154	132	54	153	175
4	112	88	74	200	189	147	66	54	76
August/ week 1	65	76	54	190	175	65	76	88	149
2	38	75	46	122	110	79	50	105	175
3	78	114	54	187	166	153	75	119	175
4	74	66	32	167	130	122	21	112	153
Sept/week 1	65	44	75	122	145	165	57	110	175
2	57	87	45	187	112	96	54	113	165
3	76	112	54	145	88	96	44	53	128
4	43	75	37	186	88	112	51	74	91
5	55	76	64	175	110	87	32	64	122

**Bacteria levels at the swimming beaches remained below the standard in 2022 with one exception in June taken at Coles Creek. Of the additional Marina bacteria samples taken, all were below the standard.*

DISCUSSION: WATER QUALITY

Water quality metrics assessed by CEMVS can be sporadic and highly variable from year to year, thus long-term data collection using consistent and comparable methodology is critical to identify trends or patterns. In general, conditions observed during 2022 did not deviate far from conditions observed during the reference period (1971-2020). Nevertheless, concerns regarding dissolved oxygen, temperature, bacteria, total phosphorus, iron, and pH were evident. In addition, CHL_a and subsequent TSI levels were indicative of a hyper eutrophic system.

In 2022 all 33 ambient discrete observations of DO were within the state guidelines except for CAR-4, Boulder Marina, Dam West Marina and Keyesport Marina. On June 23 DO was recorded at 4.13 mg/L at CAR-4 and 2.66 mg/L at Boulder Marina. On September 6 DO was 2.67 mg/L at Boulder, 4.93 mg/L at Dam West, and 3.9 at Keyesport Marinas. Since 1972, there have been 31 routine lake surface measurements observed in the summer in which DO was below 5 mg/L. DO was measured at the tail race in 15-minute intervals from April 19 through November 14, 2022. DO was recorded below the standard of 5 mg/L at the tail race during the months of May, June, July, September, and October. These lower measurements occurred during periods of low or minimal outflow, in which tainter gates were cracked or shut completely with flow coming through the sluice gate. All other measurements of DO were greater than the standard. It is not abnormal during periods of warm air and water temperatures to experience low DO. DO has an inverse relationship with temperature. As temperature increases, the ability of water to contain DO decreases, therefore the DO concentration decreases. Water temperature measurements made during 2022 indicate an increase from the historical data. This finding assumes that the historical reference 1971-2021 is the normal seasonal temperature. In a comparison of 2022 mean surface temperatures to historical mean temperatures, the water quality standard of $<2.8^{\circ}\text{C}$ was exceeded during the spring (2022 overall spring average was 6.29°C greater than the historical overall spring average). Discrete measurements of temperature were exceeded at multiple locations in the lake during the spring sample events.

E. Coliform levels were observed above the swimming standard of 235 E. Coli per 100 mL (single sample) in the lake and tributaries eleven out of the 28 samples during the four sampling events. The mean E. Coliform levels were also greater than the standard for each tributary as well as Dam West and Keyesport marinas. The highest levels were recorded in the tributaries where maximum concentrations were 825 E. Coli per 100 mL at CAR-12 and 1600 E. Coli per 100/mL at CAR-13. Bacteria levels can be highly variable and high levels may not necessarily be representative of the entire system. There were significant precipitation events which proceeded or occurred during most of the higher bacteria results recorded, though not for the highest at CAR-13 on May 17, 2022. Conversely, all of the swimming beach bacteria results monitored by project staff during the recreation season were below the standard with the exception of one (1602 E. Coli per 100 mL at Coles Creek the fourth week of June). Given that 2022 high bacteria levels in the lake and tributaries are not designated swimming areas, there is a lower risk

to humans. Long term bacteria monitoring, and analyses will be important to assess changes over time.

Phosphorus levels have surpassed the 0.05 mg/L criterion for several years. In 2022 the TP criterion was exceeded at all locations with a mean concentration across all sites of 0.26 mg/L. This is 13.4% greater than the historical mean of 0.23 mg/L, but less than the 2021 mean of 0.37 mg/L. The mean surface NO₃-N concentration in 2022 (0.50 mg/L) was less than the historic mean (1.21 mg/L) and did not exceed the criterion of 10 mg/L in 2022. Phosphorus is a limiting nutrient for primary producers (algae and plants) due to its relatively low amount in the environment. Higher inputs of TP and NO₃-N into the lake contribute to a highly productive environment which stimulates algal growth that can lead to blooms that deplete the oxygen levels during die off. In addition, blooms can sometimes contain toxins which may be harmful to humans and wildlife.

Living organisms require trace amounts of metals, excessive levels can be harmful. TFe exceeded the criterion of 1 mg/L three times at the bottom reservoir location in front of the dam (May, June, and August) and once in May at the discharge. Comparably, there are 114 times TFe was high historically (1984-2021) at the same locations. The 2022 TFe mean concentrations are comparable (RB: 1.10 mg/L, TR: 0.78) to the historical means (RB: 1.22 mg/L, TR 0.87 mg/L). Iron cycling is a function of oxidation-reduction processes. Elevated levels of iron near the bottom of a lake are not immediately detrimental to the overall lake system. Iron oxidizes relatively rapidly (minutes to hours); therefore, any iron released through the spillway will be oxidized in a short period of time.

Most freshwater streams and rivers have a natural pH ranging from 6 to 8. As pH approaches 5 (acidic), less tolerant fish and aquatic invertebrate assemblages may be extirpated, and a pH below 4.5 would be without most desired aquatic life. Conversely, when pH exceeds 9.5 (alkaline), aquatic fish and invertebrates begin to rapidly decrease and beyond 10, fish become extirpated. The pH water quality criteria for Illinois ranges from 6.5 – 9.0. The three observations pH was greater than 9.0 in 2022 were CAR-2, CAR-CSA, and CAR-KP (9.05, 9.1, 9.05 respectively). The pH means for 2022 were 8.22 (RS), 8.04 (TR), and 7.94 (TRIB) and were greater than the historical pH means.

Although there is not a state criterion for CHL_a the proposed standard of 25 mg/cm³ was exceeded at all sites in 2022. The 2022 combined CHL_a mean concentration of 41.57 mg/cm³ was greater to the historical mean of 28.74 mg/cm³. CHL_a is an indicator of the abundance of phytoplankton. Any water environment with a level recorded above 25 mg/cm³ is considered to be eutrophic (nutrient enrichment increases algal and plant growth and negative effects). The 2022 TSI level, an average of the individual trophic state indexes for Secchi depth, CHL_a, and TP, for Carlyle Lake is 75.2. Carlyle Lake is considered hyper-eutrophic based on this TSI level. This does not necessarily mean the water quality is poor, but that its trophic level indicates nutrient levels are abundant, which can support an abundance of plants and algae. Long term monitoring and analyses are important to assess changes over time.

All remaining parameters evaluated during the 2022 water quality monitoring effort were within designated criteria or within historical reference norms.

MONITORING PROGRAM RECOMMENDATIONS

The IEPA currently has listed Carlyle Lake as impaired for total phosphorous and mercury while the Kaskaskia River upstream from the Lake is impaired for fecal coliform, and mercury. In addition, the North Fork Kaskaskia River is impaired for phosphorus, Atrazine, and Terbufos. The lists of sources for these impairments are contaminated sediments, crop production, and unknown sources. At present the only tributary being sampled by CEMVS is the Kaskaskia River. IEPA also has the following listed as impaired: Hurricane Creek, North Fork Kaskaskia, and East Fork Kaskaskia. Future sampling efforts will focus on adding these three tributaries as well as mercury in the lake to the routine sampling plan to increase the dataset and improve our ability to assess the water quality condition of Carlyle Lake.

In accordance with EM-1110-2-1201, sediment samples should be taken to monitor and assess potential impacts to aquatic and human health. Sediment sampling and analyses occurred at Carlyle Lake in 2018, and prior to that in 2007. During these last analyses multiple exceedances over the recommended criteria were observed. Identifying trends over time is much more achievable with more consistent data. Contaminated sediments may have negative impacts on ecological processes. It is recommended, if possible, to sample and analyze for sediment metals and nutrients, as well as grain size analyses yearly or every two years.

Given the above-mentioned high bacteria levels observed 2022, it is recommended to continue routine bacteria sampling at all locations. This would be useful in capturing a larger picture of bacteria coming into the lake.

WORKS CITED

Carlson, R. E. (1977). A Trophic State Index for Lakes¹. *Limnology and Oceanography*, 22(2), 361-369.

USACE. (2018). *Engineering and Design: Water Quality Management*. USACE ER 1110-2-8154. Washington D.C.

USACE. (1987). *Engineering and Design: Reservoir Water Quality Analysis*. USACE ER 1110-2-1201. Washington D.C.

IEPA. (2020). <https://www2.illinois.gov/epa/topics/water-quality/watershed-management/tmdls/Pages/303d-list.aspx>

Hudson, H. (1998). Illinois Environmental Protection Agency. *Common Lake Water Quality Parameters*. Lake Notes.

APPENDIX A: FIELD DATA

Date	Location	Depth (m)	Temp (°C)	ORP (mV)	Sp Cond (µS/cm)	pH	ODO (% Sat)	ODO (mg/L)	TDS (mg/L)	Turbidity (FNU)	Secchi (in)
5/17/2022	CAR-1	0.734	20.505	326	391.1	8.32	89.4	8.04	254	19.48	
5/17/2022	CAR-12	0.663	22.382	261.7	560.3	8.27	128.1	11.1	364	26.43	
5/17/2022	CAR-13	1.097	21.743	265.8	531.3	8.02	100	8.77	345	69.83	
5/17/2022	CAR-2	1.072	22.443	250.8	358.7	9.05	143.7	12.45	233	7.73	20
5/17/2022	CAR-2	2.074	21.867	254	368.9	8.81	114.7	10.04	240	8.89	
5/17/2022	CAR-2	3.406	19.534	267.9	401.4	8.27	62.6	5.74	261	16.28	
5/17/2022	CAR-2	5.838	18.022	270.4	414.7	7.96	26.3	2.48	270	29.38	
5/17/2022	CAR-2	6.067	18.297	269.9	412.8	8.02	31.7	2.98	268	25.42	
5/17/2022	CAR-4	0.077	21.943	85	4.5	8.87	100.9	8.82	3	1.8	12
5/17/2022	CAR-4	0.978	21.946	124.8	400.3	8.68	87.2	7.63	260	44.42	
5/17/2022	CAR-4	1.91	21.615	144.7	396.5	8.79	89.9	7.91	258	45.53	
5/17/2022	CAR-4	1.934	21.684	142.9	404.4	8.47	79.3	6.96	263	53.43	
5/17/2022	CAR-4	3.043	21.553	157.4	406	8.43	73	6.43	264	64.12	
5/17/2022	CAR-4	4.2	21.459	158	404.5	8.5	72.2	6.37	263	73.48	
5/17/2022	CAR-BL-MAR	1.086	23.059	226.7	400.1	8.47	86.1	7.37	260	41.08	
5/17/2022	CAR-CSA	0.394	23.612	299	351.8	9.16	180.4	15.29	229	6.18	
5/17/2022	CAR-CSA	0.996	22.033	281.4	353.7	9.1	165.6	14.45	230	8.58	
5/17/2022	CAR-CSA	1.487	21.614	270.6	356.3	9.04	149.9	13.19	232	12.26	
5/17/2022	CAR-DW-MAR	2.13	21.67	282.3	368	8.64	100.8	8.86	239	33.31	
5/17/2022	CAR-KP-MAR	0.114	24.834	140.2	389.8	9.05	168.6	13.96	253	22.99	
5/17/2022	CAR-KP-MAR	2.601	21.78	267.3	529.6	8.12	99.7	8.75	344	70.55	
6/23/2022	CAR-1	1.615	26.142	149.8	386.6	8.06	90.6	7.32	251	15.38	
6/23/2022	CAR-12	0.323	28.036	191	430.1	8.04	140.1	10.95	280	25.91	
6/23/2022	CAR-13	1.146	26.851	187.7	485	8.01	131.9	10.53	315	27.39	
6/23/2022	CAR-13	2.097	26.88	183.9	485.3	8.03	131.1	10.45	315	30.35	
6/23/2022	CAR-2	1.16	27.97	120	374.5	8.7	176	13.77	243	8.99	19
6/23/2022	CAR-2	2.165	26.744	156.2	381	8.42	129.9	10.39	248	10.79	
6/23/2022	CAR-2	3.169	26.132	165.8	383.7	8.23	92.2	7.45	249	9.9	
6/23/2022	CAR-2	4.136	26.08	170	383.6	8.24	95.4	7.72	249	21.09	
6/23/2022	CAR-2	5.14	25.776	176.3	385.2	8.09	80.7	6.57	250	21.38	

Date	Location	Depth (m)	Temp (°C)	ORP (mV)	Sp Cond (µS/cm)	pH	ODO (% Sat)	ODO (mg/L)	TDS (mg/L)	Turbidity (FNU)	Secchi (in)
6/23/2022	CAR-2	6.112	25.521	180.1	386.7	7.94	63.7	5.21	251	23.52	
6/23/2022	CAR-4	0.433	26.537	121.7	438.4	8.24	54.4	4.36	285	41.66	13
6/23/2022	CAR-4	1.152	26.438	112.7	436.8	8.22	51.4	4.13	284	43.09	
6/23/2022	CAR-4	2.102	26.297	125.1	435.4	8.2	50.4	4.06	283	44.61	
6/23/2022	CAR-4	3.047	26.22	129.3	436.8	8.24	56.2	4.54	284	44.66	
6/23/2022	CAR-4	4.125	26.15	132.5	441.4	8.3	65.3	5.28	287	45.07	
6/23/2022	CAR-4	5.092	26.103	135.2	443.2	8.29	64.1	5.18	288	52.41	
6/23/2022	CAR-BL-MAR	0.149	29.275	179.9	389.2	8.18	85.3	6.52	253	17.58	
6/23/2022	CAR-BL-MAR	0.848	27.378	183.4	394.7	7.85	33.6	2.66	257	25.38	
6/23/2022	CAR-CSA	0.972	26.409	124.2	377.5	8.6	146.8	11.81	245	11.11	
6/23/2022	CAR-DW-MAR	0.967	26.144	141.5	383.1	8.31	101.3	8.19	249	13.91	
6/23/2022	CAR-KP-MAR	0.973	27.51	128.1	441.7	8.3	76.8	6.06	287	135.6	
8/2/2022	CAR-1	0.514	26.1	388.7	361.2	7.9	102.2	8.27	235		
8/2/2022	CAR-1	0.739	25.179	426.1	353.8	7.96	102.5	8.44	230	14.12	
8/2/2022	CAR-12	0.283	23.343	367.9	405.3	7.91	105.6	8.99	263	23.19	
8/2/2022	CAR-12	0.289	25.216	371.9	420.1	7.72	105.5	8.67	273		
8/2/2022	CAR-13	1.246	25.11	343.1	473.1	8.11	105.8	8.72	308	14.08	
8/2/2022	CAR-2	0.48	26.242	238.2	359.5	7.99	77.6	6.26	234		23
8/2/2022	CAR-2	1.214	26.26	238.7	359.5	7.99	77.6	6.26	234		
8/2/2022	CAR-2	1.219	25.378	274.8	351.9	8.1	79.2	6.49	229	18.08	
8/2/2022	CAR-2	2.416	25.325	268.4	351.9	8.08	76.7	6.29	229	20.38	
8/2/2022	CAR-2	3.21	25.253	286.7	352.1	8.06	75.7	6.22	229	20.76	
8/2/2022	CAR-2	4.138	25.238	284.6	352.2	8.05	73.9	6.07	229	20.52	
8/2/2022	CAR-2	5.362	25.221	281.6	352.3	8.04	72.3	5.94	229	23.17	
8/2/2022	CAR-2	5.982	25.222	278.9	352.3	8.03	71.9	5.91	229	24.68	
8/2/2022	CAR-4	0.535	25.669	363.2	191	7.82	82	6.69	124		13
8/2/2022	CAR-4	1.154	25.659	362.2	190.7	7.82	81.7	6.67	124		
8/2/2022	CAR-4	1.225	24.863	395.9	191.7	7.96	81.7	6.77	125	30.72	
8/2/2022	CAR-4	2.055	24.941	388.2	193	7.93	80.7	6.67	125	33.95	
8/2/2022	CAR-4	3.184	24.894	385.3	192	7.93	80.7	6.68	125	33.45	

Date	Location	Depth (m)	Temp (°C)	ORP (mV)	Sp Cond (µS/cm)	pH	ODO (% Sat)	ODO (mg/L)	TDS (mg/L)	Turbidity (FNU)	Secchi (in)
8/2/2022	CAR-4	4.066	24.79	380.3	188.8	7.9	81.3	6.75	123	33.58	
8/2/2022	CAR-4	5.113	24.784	376	190.1	7.93	81.1	6.73	124	30.25	
8/2/2022	CAR-4	6.177	24.755	373	188.3	7.93	81.1	6.73	122	34.63	
8/2/2022	CAR-BL-MAR	0.976	25.933	346.6	214.6	7.75	69	5.6	139	13.3	
8/2/2022	CAR-CSA	1.039	25.264	260.6	352.9	8.22	81.9	6.73	229	13.84	
8/2/2022	CAR-DW-MAR	1.013	25.933	298.7	345.2	8.35	91.7	7.44	224	13.41	
8/2/2022	CAR-KP-MAR	1.038	24.677	349.7	187.6	7.87	78.9	6.55	122	32.99	
9/6/2022	CAR-1	0.396	25.836	254.5	257.8	7.87	94.7	7.7	168		
9/6/2022	CAR-1	1.524	23.982	371.1	259.3	7.81	96.9	8.15	169	15.85	
9/6/2022	CAR-12	-0.144	23.233	269.3	287.9	7.16	62.7	5.35	187	9.63	
9/6/2022	CAR-12	0.165	24.77	186.8	232.2	7.25	52.5	4.36	151		
9/6/2022	CAR-13	0.692	22.552	283.3	564.5	8	107.9	9.32	367	26.3	
9/6/2022	CAR-2	0.512	26.708	171	253	8.5	113.6	9.1	164		18
9/6/2022	CAR-2	1.003	26.053	185.2	255.6	8.2	89.8	7.28	166		
9/6/2022	CAR-2	1.022	24.542	180	310.1	7.78	62.1	5.17	202		
9/6/2022	CAR-2	1.04	23.893	321.7	259.2	7.77	67.6	5.7	168	13.84	
9/6/2022	CAR-2	3.291	23.867	307.8	258.6	7.82	70.6	5.95	168	14.63	
9/6/2022	CAR-2	5.06	23.795	311.8	258.2	7.87	71.8	6.06	168	19.47	
9/6/2022	CAR-4	0.519	25.795	193.4	317.6	8.23	102.2	8.31	206		12
9/6/2022	CAR-4	1.014	22.407	287.4	322.4	7.72	64.5	5.59	210	25.53	
9/6/2022	CAR-4	1.024	24.459	183.1	311.2	7.78	62.1	5.18	202		
9/6/2022	CAR-4	3.063	22.335	214.6	322.4	7.69	62.4	5.42	210	33.4	
9/6/2022	CAR-4	5.073	22.342	241.9	325.4	7.63	54.2	4.7	211	74.7	
9/6/2022	CAR-BL-MAR	1.102	22.952	131.6	242	7.27	31.1	2.67	157	59.1	
9/6/2022	CAR-CSA	1.091	23.641	403.3	257.3	7.87	68.8	5.83	167	19.15	
9/6/2022	CAR-DW-MAR	1.017	23.856	426.3	255.3	7.74	58.4	4.93	166	20.98	
9/6/2022	CAR-KP-MAR	1.108	22.758	232.3	287.6	7.57	45.3	3.9	187	227.23	

APPENDIX B: LABORATORY DATA